

# Unit

# 7

$p$	$q$	$p \Rightarrow q$	$(p \Rightarrow q) \wedge \neg q$	$[(p \Rightarrow q) \wedge \neg q] \Rightarrow \neg p$
T	T	T	F	T
T	F	F	F	T
F	T	T	F	T
F	F	T	T	T

## MATHEMATICAL PROOFS

### Unit Outcomes:

After completing this unit, you should be able to:

- develop the knowledge of logic and logical statements.
- understand the use of quantifiers and use them properly.
- determine the validity of arguments.
- apply the principle of mathematical induction for a problem that needs to be proved inductively.
- realize the rule of inference.

### Main Contents

#### 7.1 REVISION ON LOGIC

#### 7.2 DIFFERENT TYPES OF PROOFS

#### 7.3 PRINCIPLE AND APPLICATION OF MATHEMATICAL INDUCTION

*Key Terms*

*Summary*

*Review Exercises*


## INTRODUCTION

IN ORDER TO FULLY UNDERSTAND MATHEMATICS, IT IS IMPORTANT TO UNDERSTAND WHAT A CORRECT MATHEMATICAL ARGUMENT, OR PROOF. IN THIS UNIT, YOU WILL BE INTRODUCED TO DIFFERENT METHODS OF MATHEMATICAL PROOF AND YOU WILL ALSO SEE THE ROLE OF MATHEMATICS IN PROVING MATHEMATICAL STATEMENTS. WE WILL BEGIN THE UNIT BY BRIEFLY REVISITING MATHEMATICAL LOGIC.



### OPENING PROBLEM

AFTER COMPLETING THIS UNIT, YOU SHOULD BE ABLE TO ANSWER THE FOLLOWING: CONSIDER THE FOLLOWING ARRANGEMENTS OF DOTS.



Number of dots	1	3	6	10
Sum of dots in row	1	1 + 2	1 + 2 + 3	1 + 2 + 3 + 4

NUMBERS LIKE 1, 3, 6, 10, ETC. ARE CALLED TRIANGULAR NUMBERS.

- A** CAN YOU LIST THE NEXT 5 TRIANGULAR NUMBERS?  
**B** CAN YOU GIVE A FORMULA FOR THE  $n^{\text{TH}}$  TRIANGULAR NUMBER?  
**C** LET  $T_n$  DENOTE THE  $n^{\text{TH}}$  TRIANGULAR NUMBER. CAN YOU SHOW THAT

$$\sum_{i=1}^n T_i = \frac{n(n+1)(n+2)}{6} ?$$

- D** CAN YOU FIND  $\sum_{i=1}^{40} T_i$ ?

## 7.1

### REVISION ON LOGIC

#### Revision of Statements and Logical Connectives

IN UNIT 4 OF YOUR GRADE 11 MATHEMATICS, YOU HAVE STUDIED STATEMENTS AND LOGICAL CONNECTIVES (OR OPERATORS):

NEGATION, (OR  $\vee$ ), AND ( $\wedge$ ), IMPLICATION ( $\Rightarrow$ ) AND BI-IMPLICATION ( $\Leftrightarrow$ )

THE FOLLOWING ACTIVITIES ARE DESIGNED TO HELP YOU TO REVISE THESE CONCEPTS.

## ACTIVITY 7.1



- 1 WHAT IS MEANT BY A STATEMENT (PROPOSITION)?
- 2 LIST THE PROPOSITIONAL CONNECTIVES.
- 3 WHAT IS MEANT BY A COMPOUND (COMPLEX) STATEMENT?
- 4 REVIEW THE RULES OF ASSIGNING TRUTH VALUES TO COMPOUND STATEMENTS BY COMPLETING THE TABLE BELOW WHERE  $p$  AND  $q$  ARE ANY TWO PROPOSITIONS.

$p$	$q$	$\neg p$	$p \wedge q$	$p \vee q$	$p \Rightarrow q$	$p \Leftrightarrow q$
T	T					
T	F					
F	T					
F	F					

- 5 GIVEN STATEMENTS  $p$  AND  $q$ , EACH WITH TRUTH VALUE T, FIND THE TRUTH VALUE OF EACH OF THE FOLLOWING COMPOUND STATEMENTS.

**A**  $\neg p \vee q$                       **B**  $\neg(p \vee q)$                       **C**  $\neg q \Rightarrow \neg p$

**D**  $\neg q \Leftrightarrow p$                       **E**  $\neg(p \wedge q)$


- 6 CONSTRUCT A TRUTH TABLE FOR

**A**  $\neg p \vee q$                       **B**  $(p \Rightarrow q) \Leftrightarrow \neg p$

**C**  $(p \wedge q) \Rightarrow r$                       **D**  $\neg(p \Rightarrow q) \vee \neg r$

### Open statements and quantifiers

## ACTIVITY 7.2



DECIDE WHETHER OR NOT EACH OF THE FOLLOWING IS A STATEMENT. IF IT IS A STATEMENT, DETERMINE ITS TRUTH VALUE.

- 1  $x$  IS A COMPOSITE NUMBER.
- 2 IF  $3 + 2 = 7$ , THEN  $49 = 32$ .
- 3  $x + 2 = 15$ , WHERE  $x$  IS AN INTEGER.
- 4 ALL PRIME NUMBERS ARE ODD.
- 5 THERE EXISTS A PRIME NUMBER BETWEEN 15 AND 30.
- 6 ALL BIRDS CAN FLY.

AS YOU MAY RECALL FROM YOUR LESSONS, THE WORDS **there exists** IN QUESTIONS 4, 5 AND 6 OF ACTIVITY 7.2 ARE QUANTIFIERS.

SOME OF THE SENTENCES INVOLVE VARIABLES OR UNKNOWN AND BECOME STATEMENTS. VARIABLES OR THE UNKNOWN ARE REPLACED BY SPECIFIC NUMBERS OR INDIVIDUAL. SENTENCES ARE CALLED **Statements**.

RECALL THAT OPEN STATEMENTS ARE DENOTED BY  $P(x)$  WHERE  $x$  STANDS FOR THE UNKNOWN AND STANDS FOR SOME PROPERTY THAT IS TO BE PROVED. FOR EXAMPLE, IF WE DENOTE THE OPEN STATEMENT 1) ABOVE BY  $P(x)$ , THEN  $P(x)$  STANDS FOR THE PROPERTY OF BEING A COMPOSITE NUMBER WHILE  $x$  IS THE VARIABLE OR THE UNKNOWN IN THE OPEN STATEMENT.

### Quantifiers

THERE IS A WAY OF CHANGING AN OPEN STATEMENT INTO A STATEMENT WITHOUT SUBSTITUTING INDIVIDUAL(S) FOR THE VARIABLE(S) INVOLVED BY USING WHAT WE CALL QUANTIFIERS. TWO TYPES OF QUANTIFIERS WHICH ARE USED TO CHANGE AN OPEN STATEMENT INTO A STATEMENT WITHOUT ANY SUBSTITUTION. THEY ARE:

THE UNIVERSAL QUANTIFIER DENOTED BY  $\forall$   
THE EXISTENTIAL QUANTIFIER DENOTED BY  $\exists$

THE NOTATION  $\forall$  MAY BE READ IN ANY ONE OF THE FOLLOWING WAYS:

*for all x*                      *for every x*  
*for each x*                     *for any x*

THE NOTATION  $\exists$  MAY BE READ IN ANY ONE OF THE FOLLOWING WAYS:

*there exists x,*              *for at least one x,*              *for some x*

**Example 1** LET  $P(x) \equiv x > 5$  AND  $Q(x) \equiv x$  IS AN EVEN NUMBER. THEN DETERMINE THE TRUTH VALUE OF EACH OF THE FOLLOWING STATEMENTS.

- A**  $(\forall x) P(x)$                       **B**  $(\exists x) P(x)$   
**C**  $(\exists x) [P(x) \wedge Q(x)]$         **D**  $(\forall x) [P(x) \Rightarrow Q(x)]$

**Solution**

- A**  $(\forall x) P(x)$  IS FALSE, BECAUSE IF YOU TAKE  $x = 1$ , THEN  $1 > 5$  IS FALSE.  
**B**  $(\exists x) P(x)$  IS TRUE, BECAUSE YOU CAN FIND AN  $x$  SUCH THAT  $7 > 5$  IS TRUE.  
**C**  $(\exists x) [P(x) \wedge Q(x)]$  IS TRUE, IF YOU TAKE  $x = 6$ , THEN  $6 > 5$  AND 6 IS EVEN.  
**D**  $(\forall x) [P(x) \Rightarrow Q(x)]$  IS FALSE, FOR,  $P(7)$  IS TRUE BUT  $Q(7)$  IS FALSE.

**Example 2** CHANGE THE FOLLOWING OPEN STATEMENT INTO A STATEMENT AND DETERMINE THE TRUTH VALUE.

$P(x): x^2 < 0$ , WHERE  $x$  IS A COMPLEX NUMBER.

**Solution**

USING THE UNIVERSAL QUANTIFIER,  $(\forall x) P(x)$  IS FALSE, BECAUSE WHEN  $x$  IS A REAL NUMBER SUCH AS  $x = 1$ ,  $1^2 < 0$  IS FALSE.

USING THE EXISTENTIAL QUANTIFIER,  $(\exists x) P(x)$  IS TRUE BECAUSE WHEN  $x$  IS AN IMAGINARY NUMBER SUCH AS  $x = 2i$ , ETC,  $(2i)^2 = -4$ , ETC.

**Exercise 7.1**

- 1 LET  $P(x) = x$  IS A STUDENT WHO STUDIED GEOMETRY.  
 THEN,  $\forall x P(x)$  IS READ AS: \_\_\_\_\_  
 WHILE  $\exists x P(x)$  IS READ AS: \_\_\_\_\_
- 2 GIVEN THE OPEN STATEMENTS:  
 $P(x) \equiv x$  IS A PRIME NUMBER.  $Q(x) \equiv x$  IS AN ODD NUMBER.  
 DETERMINE THE TRUTH VALUE OF EACH OF THE FOLLOWING STATEMENTS.

**A**  $(\forall x) P(x)$                       **B**  $(\exists x) P(x)$                       **C**  $(\exists x) (\neg P(x))$   
**D**  $(\forall x) [P(x) \Rightarrow Q(x)]$       **E**  $(\exists x) [P(x) \wedge \neg Q(x)]$
- 3 IF  $x$  AND  $y$  ARE INTEGERS, DETERMINE THE TRUTH VALUE OF EACH OF THE FOLLOWING.

**A**  $(\exists x) (\forall y) (x \leq y)$                       **B**  $(\exists x) (\forall y) (x^2 \leq y)$   
**C**  $(\forall x) (\exists y) (x \leq y)$                       **D**  $(\forall x) (\forall y) (x + y = y + x)$   
**E**  $(\forall x) (\exists y) (x + y = 0)$
- 4 EXPRESS EACH OF THE FOLLOWING USING QUANTIFIERS.

**A** SOME STUDENTS IN THIS CLASS HAVE VISITED GONDAR.  
**B** EVERY STUDENT IN THIS CLASS HAS VISITED EITHER GONDAR OR

**Arguments and validity of arguments**

**ACTIVITY 7.3**



- 1 DISCUSS WHETHER OR NOT THE FOLLOWING ARGUMENTS ARE VALID.

**A** IF THE DAY IS CLOUDY, THEN IT RAINS.  
 DOES THIS MEAN THAT IF IT RAINS, THERE ARE CLOUDS?

**B** IF  $x$  IS A PRIME NUMBER AND COMPOSITE NUMBER, THEN  $x$  IS A COMPOSITE NUMBER.
- 2 CONSTRUCT A SINGLE TRUTH TABLE FOR THE FOLLOWING STATEMENTS  $p \Rightarrow q, \neg q \Rightarrow r$ , AND  $p \Rightarrow r$ .  
 FIND OUT THE ROWS IN WHICH THE STATEMENTS  $p \Rightarrow q$  AND  $\neg q \Rightarrow r$  ARE BOTH TRUE BUT  $p \Rightarrow r$  IS FALSE.

AN ARGUMENT IS AN ASSERTION THAT A GIVEN SET OF STATEMENTS CALLED THE (hypothesis), YIELDS ANOTHER STATEMENT, CALLED (the consequent).

AN ARGUMENT IS SAID TO BE VALID, IF AND ONLY IF THE CONJUNCTION OF ALL THE PREMISES IMPLIES THE CONCLUSION. IN OTHER WORDS, IF WE ASSUME THAT THE STATEMENTS IN THE PREMISES ARE ALL TRUE, THEN (FOR A VALID ARGUMENT), THE CONCLUSION MUST BE TRUE. AN ARGUMENT WHICH IS NOT VALID IS CALLED A FALLACY.

THE VALIDITY OF AN ARGUMENT CAN EASILY BE CHECKED BY CONSTRUCTING A TRUTH TABLE. YOU MUST SHOW THAT THE PREMISES ALTOGETHER ALWAYS IMPLY THE CONCLUSION. IN OTHER WORDS, YOU SHOW THAT "CONJUNCTION OF THE PREMISES" IS ALWAYS TRUE (OR A TAUTOLOGY).

TO SHOW THE VALIDITY OF AN ARGUMENT, YOU HAVE TO SHOW THAT THE CONCLUSION IS ALWAYS TRUE WHENEVER ALL THE PREMISES ARE TRUE.

**Example 3** IS THE FOLLOWING ARGUMENT VALID?

IF I AM RICH, THEN I AM HEALTHY.  
 I AM HEALTHY.  
 THEREFORE, I AM RICH.

**Solution**

NOTE THAT THE FIRST TWO STATEMENTS ARE THE PREMISES WHILE THE LAST STATEMENT IS THE CONCLUSION. THIS ARGUMENT IS NOT A VALID ARGUMENT. TO SEE WHY, WE SHALL FIRST SYMBOLIZE IT.

LET  $p$  STAND FOR THE STATEMENT "I AM RICH" AND  $q$  FOR THE STATEMENT "I AM HEALTHY".

THEN, THE SYMBOLIC FORM OF THE ABOVE ARGUMENT BECOMES:

$$p \Rightarrow q$$

$$\frac{q}{p} \quad \text{OR} \quad p \Rightarrow q, q \vdash p$$

THIS ARGUMENT WOULD BE VALID, IF THE IMPLICATION WERE ALWAYS TRUE.

WHEN YOU CONSTRUCT THE TRUTH TABLE FOR THIS CONDITIONAL STATEMENT AS SHOWN IN THE TABLE, YOU WILL SEE THAT THE CONCLUSION COULD BE FALSE WHILE BOTH THE PREMISES ARE TRUE. (SEE THE THIRD COLUMN). IN OTHER WORDS,  $(p \Rightarrow q) \wedge q \Rightarrow p$  IS NOT A TAUTOLOGY. THUS, THE ARGUMENT IS INVALID.

$p$	$q$	$p \Rightarrow q$	$(p \Rightarrow q) \wedge q$	$[(p \Rightarrow q) \wedge q] \Rightarrow p$
T	T	T	T	T
T	F	F	F	T
F	T	T	T	F
F	F	T	F	T

**Example 4** IS THE FOLLOWING ARGUMENT VALID?

IF I AM HEALTHY, THEN I WILL BE HAPPY.  
 I AM NOT HAPPY.  
 THEREFORE, I AM NOT HEALTHY.

**Solution** ONCE AGAIN, TO CHECK THE VALIDITY OF THIS ARGUMENT, LET  $p$  REPRESENT "I AM HEALTHY" AND  $q$  REPRESENT "I AM HAPPY". THE SYMBOLIC FORM OF THE ARGUMENT IS:

$$\begin{array}{l}
 p \Rightarrow q \\
 \frac{\neg q}{\neg p} \\
 p \Rightarrow q, \neg q \vdash \neg p.
 \end{array}$$

THIS ARGUMENT WILL BE VALID, IF THE IMPLICATION  $[(p \Rightarrow q) \wedge \neg q] \Rightarrow \neg p$  IS ALWAYS TRUE (A TAUTOLOGY). CONSTRUCTING A TRUTH TABLE AS SHOWN BELOW, YOU NOTICE ARGUMENT IS VALID.

$p$	$q$	$\neg p$	$\neg q$	$p \Rightarrow q$	$(p \Rightarrow q) \wedge \neg q$	$[(p \Rightarrow q) \wedge \neg q] \Rightarrow \neg p$
T	T	F	F	T	F	T
T	F	F	T	F	F	T
F	T	T	F	T	F	T
F	F	T	T	T	T	T

**Example 5** SHOW THAT THE FOLLOWING ARGUMENT IS VALID.

IF YOU SEND ME AN EMAIL, THEN I WILL FINISH WRITING MY PROJECT.  
 IF I FINISH WRITING MY PROJECT, THEN I WILL GET RELAXED.  
 THEREFORE, IF YOU SEND ME AN EMAIL, THEN I WILL GET RELAXED.

**Solution**

LET:  $p \equiv$  YOU SEND ME AN EMAIL  
 $q \equiv$  I FINISH WRITING MY PROJECT  
 $r \equiv$  I GET RELAXED. THEN THE SYMBOLIC FORM OF THIS ARGUMENT WILL BE AS FOLLOWS:

$$\begin{array}{l}
 p \Rightarrow q \\
 \frac{q \Rightarrow r}{p \Rightarrow r}
 \end{array}$$

NOW, THE IMPLICATION  $[(p \Rightarrow q) \wedge (q \Rightarrow r)] \Rightarrow (p \Rightarrow r)$  IS ALWAYS TRUE AS SHOWN IN THE TRUTH TABLE BELOW.

$p$	$q$	$r$	$p \Rightarrow q$	$q \Rightarrow r$	$p \Rightarrow r$	$(p \Rightarrow q) \wedge (q \Rightarrow r)$	$[(p \Rightarrow q) \wedge (q \Rightarrow r)] \Rightarrow (p \Rightarrow r)$
T	T	T	T	T	T	T	T
T	T	F	T	F	F	F	T
T	F	T	F	T	T	F	T
T	F	F	F	T	F	F	T
F	T	T	T	T	T	T	T
F	T	F	T	F	T	F	T
F	F	T	T	T	T	T	T
F	F	F	T	T	T	T	T

THEREFORE, THE ARGUMENT  $p \Rightarrow r \vdash p \Rightarrow r$  IS VALID.

THE CONSTRUCTION OF SUCH A BIG TRUTH TABLE MAY BE AVOIDED BY STUDYING AND APPLYING THE FOLLOWING RULES BY WHICH WE CHECK WHETHER A GIVEN ARGUMENT IS NOT. THEY ARE CALLED **inference** AND ARE LISTED AS FOLLOWS.

<b>1</b>	$\frac{p}{p \vee q}$	<b>PRINCIPLE OF ADJUNCTION.</b> IT STATES THAT IF <b>IS TRUE, THEN IS ALSO TRUE FOR ANY PROPOSITION</b>
<b>2</b>	$\frac{p \wedge q}{p}$	<b>PRINCIPLE OF DETACHMENT.</b> IT STATES THAT IF <b>IS TRUE, THEN IS TRUE".</b>
<b>3</b>	$\frac{p}{p \wedge q}$	<b>PRINCIPLE OF CONJUNCTION.</b> IT STATES THAT WHENEVER <b>ARE TRUE THE STATEMENT IS ALSO TRUE.</b>
<b>4</b>	$\frac{p}{q}$	<b>Modus ponens.</b> IT STATES THAT WHENEVER THE <b>IMPLICATION AND THE HYPOTHESIS IS TRUE, THEN THE CONSEQUENT IS TRUE. RECALL THE RULE OF IMPLICATION.</b>
<b>5</b>	$\frac{\neg q}{\neg p}$	<b>Modus Tollens.</b> IT STATES THAT WHENEVER <b>IS TRUE AND IS FALSE, THEN IS ALSO FALSE.</b>
<b>6</b>	$\frac{q \Rightarrow r}{p \Rightarrow r}$	<b>PRINCIPLE OF TOLLENS (LAW OF SYLLOGISM).</b> IT MAY BE REMEMBERED AS THE TRANSITIVE PROPERTY OF IMPLICATION. THIS LAW WAS ONE OF ARISTO (384 – 322 B.C.) MAIN CONTRIBUTIONS TO LOGIC.
<b>7</b>	$\frac{\neg p}{q}$	<b>Modus Tollens Ponens.</b> THIS RULE IS ALSO CALLED <b>THE syllogism.</b>

LET US NOW CONSIDER EXAMPLES THAT SHOW HOW THE ABOVE RULES OF INFERENCE ARE APPLIED FOR FINDING OF T

**Example 6** IDENTIFY THE RULE OF INFERENCE APPLIED FOR FINDING OF T ARGUMENTS.

**A** IT IS RAINING.

THEREFORE, IT IS RAINING OR IT IS COLD.

*The rule that applies to this argument is rule 1 (adjunction).*



- B** ABDISSA IS RICH AND HAPPY.  
THEREFORE, HE IS RICH.  
*The rule applied here is rule 2 (Detachment).*
- C** IT IS COLD TODAY.  
IT IS RAINING TODAY.  
THEREFORE, IT IS RAINING AND IT IS COLD TODAY.  
*This argument uses rule 3 (conjunction).*
- D** IF HANNA WORKS HARD, THEN SHE WILL SCORE GOOD GRADES.  
THEREFORE HANNA SCORES GOOD GRADES.  
*This argument uses rule 4 (Modus ponens).*
- E** IF IT IS RAINING, THEN I GET WET WHEN I GO OUTSIDE.  
IT IS NOT RAINING.  
THEREFORE, IT IS NOT RAINING.  
*In this argument, the appropriate rule is rule 5 (Modus Tollens).*
- F** IF I GET A JOB, THEN I WILL EARN MONEY.  
IF I EARN MONEY, THEN I WILL BUY A COMPUTER.  
THEREFORE, IF I GET A JOB, THEN I WILL BUY A COMPUTER.  
*The inference rule 6 (Principle of syllogism) is applied here.*
- G** EITHER WAGES ARE LOW OR PRICES ARE HIGH.  
WAGES ARE NOT LOW.  
THEREFORE, PRICES ARE HIGH.  
*The inference rule applied here is rule 7. (Modus Tollens Ponens)*

**Example 7** USING RULES OF INFERENCE, CHECK THE VALIDITY OF ARGUMENT.

$$\begin{array}{l} p \\ p \Rightarrow q \\ \frac{q \Rightarrow r}{r} \end{array}$$

**Solution**

- 1**  $p$  IS TRUE (PREMISE)
  - 2**  $p \Rightarrow q$  IS TRUE (PREMISE)
  - 3**  $q$  IS TRUE (MODUS PONENS FROM 1, 2)
  - 4**  $q \Rightarrow r$  IS TRUE (PREMISE)
  - 5**  $r$  IS TRUE (MODUS PONENS FROM 3, 4)
- THEREFORE, THE ARGUMENT IS VALID, I.E.,  $r \vdash r$  IS VALID.

**Note:**

THIS IS NOT THE ONLY WAY YOU CAN SHOW THIS. HERE IS ANOTHER SET OF STEPS.

- 1  $p$  IS TRUE (PREMISE).
  - 2  $p \Rightarrow q$  IS TRUE (PREMISE).
  - 3  $q \Rightarrow r$  IS TRUE (PREMISE).
  - 4  $p \Rightarrow r$  IS TRUE (SYLLOGISM FROM 2,3).
  - 5  $r$  IS TRUE (MODUS PONENS FROM 1, 4).
- THEREFORE, THE ARGUMENT IS VALID.

ALL THE EXAMPLES CONSIDERED ABOVE ARE EXAMPLES OF VALID ARGUMENTS. IT IS NOW SEE AN EXAMPLE OF AN INVALID ARGUMENT (OR A FALLACY).

**Example 8** 
$$\frac{q \quad \neg p \Rightarrow \neg q}{\neg p}$$

**Solution**

- 1  $q$  IS TRUE (PREMISE)
  - 2  $\neg q$  IS FALSE FROM (1)
  - 3  $\neg p \Rightarrow \neg q$  IS TRUE (PREMISE)
  - 4  $\neg p$  IS FALSE (FROM 2 AND 3)
- THEREFORE, THE ARGUMENT FORM IS NOT VALID.

**Exercise 7.2**

- 1 WHICH OF THE FOLLOWING ARE STATEMENTS AND WHICH OF THEM ARE NOT STATEMENTS?
  - A PLATO WAS A PHILOSOPHER.
  - B  $\sqrt{3}$  IS RATIONAL
  - C  $x^2 + 1 = 5$
  - D  $(\exists x)(x^2 + 1 = 5)$
  - E WHAT IS TODAY'S DATE?
- 2 LET  $p: 5 + 3 = 9$  AND  $q: \text{TODAY IS SUNNY}$ 
  - A WRITE EACH OF THE FOLLOWING IN SYMBOLIC FORM
    - I  $5 + 3 = 9$  OR TODAY IS NOT SUNNY
    - II  $5 + 3 = 9$  ONLY IF TODAY IS SUNNY
    - III  $5 + 3 \neq 9$  IF AND ONLY IF TODAY IS SUNNY
    - IV IT IS SUFFICIENT THAT TODAY IS SUNNY IN ORDER THAT
  - B WRITE EACH OF THE FOLLOWING IN WORDS.
    - I  $p \wedge \neg q$
    - II  $\neg p \Rightarrow q$
    - III  $(p \vee q) \Rightarrow \neg q$

3 USING TRUTH TABLES, SHOW THAT EACH PAIR OF STATEMENTS ARE EQUIVALENT.

**A**  $\neg p \vee q ; \neg q \Rightarrow \neg p$                       **B**  $\neg p \Leftrightarrow \neg q ; p \Leftrightarrow q$

**C**  $\neg p \Leftrightarrow q ; (p \vee q) \wedge \neg(p \wedge q)$

4 USING TRUTH TABLES, CHECK WHETHER EACH ONE OF ARGUMENTS GIVEN SYMBOLICALLY IS VALID OR INVALID (A FALLACY).

**A**  $\frac{p}{q}$                       **B**  $\frac{p}{\neg q}$                       **C**  $\frac{p \wedge q}{p \Rightarrow q}$   
 $\frac{q}{p \Rightarrow q}$                        $\frac{\neg q}{q \Rightarrow p}$                        $\frac{p \Rightarrow q}{p \vee q}$

5 FOR EACH OF THE FOLLOWING ARGUMENTS ~~SWITCH THEM~~ DETERMINE WHETHER THE ARGUMENT IS VALID OR NOT.

**A** YOUR TROUBLES START WHEN YOU GET MARRIED.  
 YOU HAVE NO TROUBLES.  
 THEREFORE, YOU ARE NOT MARRIED.

**B** IF LEGESSE DRINKS BEER, HE IS AT LEAST 18 YEARS OLD  
 LEGESSE DOES NOT DRINK BEER.  
 THEREFORE, LEGESSE IS NOT YET 18 YEARS OLD.

6 USING RULES OF INFERENCE CHECK THE VALIDITY OF EACH ARGUMENTS.

$p \Rightarrow (q \vee r)$   
**A**  $\frac{\neg q \wedge \neg r}{\neg p}$

**B** IF I STUDY, THEN I WILL NOT FAIL IN MATHEMATICS.  
 IF I DO NOT WATCH TV FREQUENTLY, THEN I WILL STUDY. BUT, I FAILED IN MATHEMATICS.  
 THEREFORE, I MUST HAVE WATCHED TV FREQUENTLY.

7 USING TRUTH TABLES, CHECK THE VALIDITY OF EACH ARGUMENTS.

$p \wedge \neg q$                        $p \wedge \neg q$                        $p \vee \neg r$   
**A**  $\frac{r \vee \neg p}{q \Rightarrow r}$                       **B**  $\frac{\neg q}{p}$                       **C**  $\frac{p \vee q}{q \vee r}$

8 USING RULES OF INFERENCE CHECK THE VALIDITY OF EACH ARGUMENTS.

**A**  $p \Rightarrow \neg q, r \Rightarrow q, r \vdash \neg p$

**B** HAILU'S BOOKS ARE ON THE DESK OR ON THE SHELF.  
 THE BOOKS ARE NOT ON THE SHELF.  
 THEREFORE, THEY ARE ON THE DESK

**C** IF 5 IS EVEN, THEN 2 IS PRIME. 2 IS PRIME AND IS POSITIVE.  
 4 IS NOT POSITIVE.  
 THEREFORE, 5 IS NOT EVEN.

## 7.2 DIFFERENT TYPES OF PROOFS


IN MATHEMATICS, A PROOF OF A GIVEN STATEMENT IS A SEQUENCE OF STATEMENTS THAT ARGUMENT. WHEN A VALID ARGUMENT IS CONSTRUCTED, YOU SAY THAT THE GIVEN STATEMENT IS PROVED. THERE ARE DIFFERENT METHODS BY WHICH PROOFS ARE CONSTRUCTED. THE METHODS OF INFERENCE DISCUSSED ABOVE, ARE INSTRUMENTS TO CONSTRUCT PROOFS. IN THIS SECTION, WE WILL CONSIDER DIFFERENT TYPES OF PROOFS OF MATHEMATICAL STATEMENTS.

SINCE MANY MATHEMATICAL STATEMENTS ARE IMPLICATIONS, THE TECHNIQUES FOR PROVING IMPLICATIONS ARE IMPORTANT. RECALL THAT IF  $p$  IS TRUE AND  $q$  IS FALSE, THEN  $p \Rightarrow q$  IS FALSE. THEREFORE, YOU NOTICE THAT WHEN  $p$  IS TRUE, THE ONLY THING TO BE SHOWN IS THAT  $q$  IS TRUE; IT IS NOT USUALLY THE CASE THAT  $q$  IS TRUE, IN ISOLATION. THE FOLLOWING DISCUSSION WILL GIVE YOU THE MOST COMMON TECHNIQUES FOR PROVING IMPLICATIONS.

### Direct proof

THE IMPLICATION  $p \Rightarrow q$  CAN BE PROVED BY SHOWING THAT IF  $p$  IS TRUE, THEN  $q$  MUST ALSO BE TRUE. A PROOF OF THIS KIND IS CALLED a **direct proof**. TO CONSTRUCT SUCH A PROOF, YOU ASSUME THAT  $p$  IS TRUE AND USE RULES OF INFERENCE AND FACTS ALREADY KNOWN OR PROVEN TO SHOW THAT  $q$  MUST ALSO BE TRUE.

### ACTIVITY 7.4



- 1 COMPLETE THE PROOF OF THE FOLLOWING STATEMENT: IF  $x$  AND  $y$  ARE ODD INTEGERS, THEN  $x + y$  IS AN EVEN INTEGER.

**Proof:**

IF  $x$  AND  $y$  ARE ODD INTEGERS, THEN THERE EXIST INTEGERS  $n$  AND  $m$  SUCH THAT

$$x = 2n + 1 \text{ AND } y = 2m + 1.$$

$$\Rightarrow x + y = \underline{\hspace{2cm}}$$

⋮

THEREFORE,  $x + y$  IS AN EVEN INTEGER.

- 2 GIVEN BELOW IS A PROOF OF THE FOLLOWING STATEMENT: IF EACH OF THE STATEMENTS IN THE PROOF IS TRUE, THEN  $\frac{m+5}{n+5} > \frac{m}{n}$ . INVESTIGATE WHY EACH OF THE STATEMENTS IN THE PROOF IS TRUE.

$$\forall n, m \in \mathbb{R}, \text{ IF } n > m > 0, \text{ THEN } \frac{m+5}{n+5} > \frac{m}{n}.$$

**Proof:**

$$n > m \Rightarrow 5n > 5m \Rightarrow 5n + mn > 5m + mn \Rightarrow n(m+5) > m(n+5)$$

$$\Rightarrow \frac{m+5}{n+5} > \frac{m}{n}$$

**Example 1** GIVE A DIRECT PROOF OF THE STATEMENT, "IF  $n$  IS ODD, THEN  $n^2 + 1$  IS ODD".

**Proof:**

ASSUME THAT THE HYPOTHESIS OF THE STATEMENT IS TRUE, I.E. SUPPOSE THAT  $n$  IS ODD. THEN  $n = 2k + 1$  FOR SOME INTEGER

THEN, IT FOLLOWS THAT  $(n^2 + 1) = (2k + 1)^2 + 1 = 4k^2 + 4k + 1 + 1 = 4k^2 + 4k + 2 = 2(2k^2 + 2k + 1) = 2m + 2$  (WHERE  $m = 2k^2 + 2k$  WHICH IS AN INTEGER).

THEREFORE,  $n^2 + 1$  IS ODD (AS IT IS 1 MORE THAN AN EVEN INTEGER).

### The method of cases or exhaustion

IN THIS METHOD, EACH AND EVERY POSSIBLE CASE IS CONSIDERED.

**Example 2** SHOW THAT  $3n + 7$  IS ODD FOR ALL  $n \in \mathbb{Z}$

**Proof:**

**Case 1**  $n$  IS EVEN

$n$  IS EVEN  $\Rightarrow n = 2k$ , FOR  $k \in \mathbb{Z}$ , BY DEFINITION.

$$\Rightarrow n^2 + 3n + 7 = (2k)^2 + 3(2k) + 7 = 4k^2 + 6k + 7 = 2(2k^2 + 3k + 3) + 1$$

HENCE  $n^2 + 3n + 7$  IS ODD.

**Case 2**  $n$  IS ODD

$n$  IS ODD  $\Rightarrow n = 2k + 1$ , FOR SOME  $k \in \mathbb{Z}$

$$\begin{aligned} \text{ACCORDINGLY } n^2 + 3n + 7 &= (2k + 1)^2 + 3(2k + 1) + 7 = 4k^2 + 4k + 1 + 6k + 3 + 7 \\ &= 4k^2 + 10k + 11 = 2(2k^2 + 5k + 5) + 1 \end{aligned}$$

THUS  $n^2 + 3n + 7$  IS ODD

$\therefore$  FROM CASES 1 AND 2,  $n^2 + 3n + 7$  IS ODD  $\forall n \in \mathbb{Z}$ .

**Example 3** SHOW THAT FOR ANY  $x, y$ , THE MAXIMUM OF  $\frac{x + y + |x - y|}{2}$  IS GIVEN BY

$$\frac{x + y + |x - y|}{2}$$

**Proof:**

TWO CASES ARISE: EITHER  $x < y$

**Case 1**  $x \geq y$

$$x \geq y \Rightarrow x - y \geq 0$$

THEN THE MAXIMUM OF  $\frac{x + y + |x - y|}{2}$  IS  $x$  AND  $|x - y| = x - y$  BY DEFINITION OF ABSOLUTE VALUE.

$$\text{NOW, } \frac{x + y + |x - y|}{2} = \frac{x + y + (x - y)}{2} = \frac{2x}{2} = x$$

$$\text{HENCE THE MAXIMUM OF } \frac{x + y + |x - y|}{2} \text{ IS } x$$

Case 2  $x < y$

$$x < y \Rightarrow x - y < 0 \Rightarrow \text{MAXIMUM OF } x \text{ AND } y \text{ IS } y \text{ AND } |x - y| = -(x - y) = -x + y.$$

$$\text{HERE, } \frac{x + y + |x - y|}{2} = \frac{x + y - (x - y)}{2} = \frac{2y}{2} = y$$

$$\text{SO THE MAXIMUM OF } x \text{ AND } y \text{ IS } \frac{x + y + |x - y|}{2} = y$$

$$\therefore \text{ THE MAXIMUM OF } x \text{ AND } y \text{ IS OR GIVEN BY } \frac{x + y + |x - y|}{2}.$$

### Indirect proof

SINCE THE IMPLICATIONS EQUIVALENT TO ITS CONTRAPOSITIVE IMPLICATION  $p \Rightarrow q$  CAN BE PROVED BY PROVING ITS CONTRAPOSITIVE, TRUE STATEMENT. A PROOF THAT USES THIS TECHNIQUE IS CALLED AN

**Example 4** PROVE THE STATEMENT "IF  $n$  IS ODD, THEN  $n^2$  IS ODD".

**Proof:**

ASSUME THAT THE CONCLUSION OF ONE IS FALSE, I.E. SUPPOSE  $n$  IS EVEN. THEN  $n = 2k$  FOR SOME INTEGER  $k$ . IT FOLLOWS THAT

$$n^2 = (2k)^2 = 4k^2 = 2(2k^2).$$

SO  $n^2$  IS EVEN (AS IT IS A MULTIPLE OF 2).

THUS, YOU HAVE SHOWN THAT IF  $n$  IS EVEN, THEN  $n^2$  IS EVEN. YOU SHOWED THAT THE NEGATION OF THE CONCLUSION IMPLIES THE NEGATION OF THE HYPOTHESIS. THEREFORE, THE CONTRAPOSITIVE, WHICH SAYS "IF  $n^2$  IS ODD, THEN  $n$  IS ODD" IS TRUE.

THIS ENDS THE PROOF.

#### Remark:

IN EXAMPLE 4 THE STATEMENT "IF  $n$  IS ODD, THEN  $n^2$  IS ODD" IS PROVED. USING THE METHOD OF EXAMPLE 5 WE HAVE EQUALLY PROVED THAT THE STATEMENT "IF  $n^2$  IS EVEN, THEN  $n$  IS EVEN" IS ALSO TRUE, BECAUSE THIS STATEMENT IS THE CONTRAPOSITIVE OF THE ABOVE ONE.

**Example 5** SHOW THAT IF  $x, y \in \mathbb{R}$ , WITH  $x > 0$  AND  $y > 0$ , THEN

$$\text{IF } xy > 25 \text{ THEN } x > 5 \text{ OR } y > 5.$$

**Proof:**

YOU CAN USE INDIRECT PROOF.

SUPPOSE,  $0 < x \leq 5$  AND  $0 < y \leq 5$ . THEN,  $0 < xy \leq 5(5)$ . I.E.,  $0 < xy \leq 25$ .

THUS, THE PRODUCT IS NOT LARGER THAN 25.

$\therefore$  IF  $xy > 25$ , THEN  $x > 5$  OR  $y > 5$  BY A CONTRA POSITIVE.

## Proof by contradiction

IN THE PREVIOUS METHODS OF PROOF, YOU USE PROOF BY CONTRADICTION TO ASSUME THE STATEMENT IS FALSE AND FINALLY CONCLUDE IT IS TRUE. NOW WHAT WILL HAPPEN IF YOU START BY ASSUMING THE STATEMENT IS TRUE AND FINALLY CONCLUDE IT IS FALSE? IF THIS ASSUMPTION LEADS TO A CONCLUSION WHICH CONTRADICTS EITHER ONE OF THE ASSUMPTIONS OR A PREVIOUSLY KNOWN FACT, THEN THE ASSUMPTION WAS NOT CORRECT. THIS WILL TELL YOU THAT THE STATEMENT IS ALWAYS TRUE. THIS METHOD OF ARGUMENT IS KNOWN AS PROOF BY CONTRADICTION.

**Example 6** PROVE THE FOLLOWING STATEMENT BY USING THE METHOD OF CONTRADICTION: "2 IS AN IRRATIONAL NUMBER".

**Proof:**

LET  $p$  BE THE STATEMENT "2 IS AN IRRATIONAL NUMBER". SUPPOSE THAT  $p$  IS FALSE, THEN  $\sqrt{2}$  IS A RATIONAL NUMBER. WE SHALL NOW SHOW THAT THIS LEADS TO A CONTRADICTION. THE ASSUMPTION THAT  $\sqrt{2}$  IS RATIONAL IMPLIES THAT THERE EXIST INTEGERS

$a$  AND  $b$  SUCH THAT  $\sqrt{2} = \frac{a}{b}$ , WHERE  $a$  AND  $b$  HAVE NO COMMON FACTOR OTHER THAN 1 (SO THAT  $\frac{a}{b}$  IS IN ITS LOWEST TERMS).

SINCE  $\sqrt{2} = \frac{a}{b}$ , BY SQUARING BOTH SIDES YOU GET

$$2 = \frac{a^2}{b^2} \Rightarrow a^2 = 2b^2.$$

THIS MEANS THAT  $a^2$  IS EVEN IMPLYING  $a$  IS EVEN. NOW, SINCE  $a^2 = 2b^2$ , IT FOLLOWS THAT  $a^2 = 2c$  FOR SOME INTEGER  $c$ .

THUS,  $a^2 = 2c \Rightarrow a^2 = 4c^2$ .

THIS AGAIN MEANS THAT  $a^2$  IS EVEN, HENCE  $a$  IS EVEN AS WELL. HENCE 2 IS A COMMON FACTOR OF  $a$  AND  $b$ .

NOTICE THAT IT HAS BEEN SHOWN THAT  $\sqrt{2}$  IS RATIONAL. NOTE THAT AS SHOWN

ABOVE, FROM  $\sqrt{2} = \frac{a}{b}$  IS RATIONAL AND  $a$  AND  $b$  HAVE NO COMMON FACTOR OTHER THAN 1

AND AT THE SAME TIME 2 DIVIDES BOTH  $a$  AND  $b$ . THIS IS A CONTRADICTION.

THIS IS A CONTRADICTION, SINCE YOU HAVE SHOWN THAT  $\sqrt{2}$  IS RATIONAL WHERE

$a$  AND  $b$  ARE INTEGERS WITH NO COMMON FACTOR OTHER THAN 1.

HENCE  $p$  IS FALSE, AS A RESULT, "2 IS AN IRRATIONAL NUMBER" IS TRUE.

**Example 7** SHOW THAT THE SUM OF A RATIONAL AND AN IRRATIONAL NUMBER IS AN IRRATIONAL NUMBER.

**Proof:**

LET  $r$  BE A RATIONAL NUMBER AND  $s$  BE AN IRRATIONAL NUMBER.

SUPPOSE THAT ON THE CONTRARY,  $r + s$  IS RATIONAL.

THEN  $a = \frac{p}{q}$  AND  $b = \frac{r}{s}$  FOR SOME  $p, r, s \in \mathbb{Z}, q, s \neq 0$ .

NOW,  $a + b = \frac{p}{q} + \frac{r}{s} = \frac{ps + qr}{sq}$   
 $\Rightarrow a + b$  IS RATIONAL.  $ps + qr \in \mathbb{Z}$  AND  $sq \in \mathbb{Z}, sq \neq 0$

THIS CONTRADICTS THE ASSUMPTION THAT  $a + b$  IS IRRATIONAL.

THUS,  $\sqrt{2}$  IS RATIONAL. ~~IS IRRATIONAL~~

### Disproving by counter-example

## ACTIVITY 7.5



GIVE THE NEGATION OF EACH OF THE FOLLOWING STATEMENTS IN SYMBOLIC FORM.

- 1  $(\forall x) (x^2 > 0)$ , WHERE  $x$  IS A REAL NUMBER
- 2  $(\exists x) (2x \text{ IS A PRIME NUMBER})$ , WHERE  $x$  IS A NATURAL NUMBER
- 3  $((\forall x) (\exists y) (x = y^2 + 1))$ , WHERE  $x$  AND  $y$  ARE REAL NUMBERS

#### Note:

FROM ACTIVITY 7.5, YOU HAVE THE FOLLOWING RESULTS:

- 1  $\neg(\forall x) (P(x)) = (\exists x) (\neg P(x))$
- 2  $\neg(\exists x) (Q(x)) = (\forall x) (\neg Q(x))$

SUPPOSE THAT YOU WANT TO SHOW THAT A STATEMENT IS FALSE. THIS IS DONE BY PRODUCING AN ELEMENT FROM THE UNIVERSAL SET THAT WHEN SUBSTITUTED IN PLACE OF AN ELEMENT IS CALLED A COUNTEREXAMPLE.

NOTE THAT ONLY ONE COUNTEREXAMPLE NEEDS TO BE FOUND TO SHOW THAT (

**Example 8** DISPROVE THE STATEMENT:

"FOR EVERY NATURAL NUMBER  $n$ ,  $n^2 + 121$  IS PRIME"

**Proof:**

IT IS SUFFICIENT TO FIND ONE NATURAL NUMBER THAT SATISFIES THIS CONDITION. THUS, IF YOU TAKE  $n = 5$ , YOU SEE THAT  $5^2 + 121 = 91$ . BUT 91 IS NOT A PRIME NUMBER AS 7 DIVIDES 91 I.E.  $91 = 7 \times 13$ .

THEREFORE, THE STATEMENT " $n^2 + 121$  IS PRIME" IS NOW DISPROVED USING THE COUNTEREXAMPLE.

THE DIFFERENT METHODS OF PROOFS DISCUSSED ABOVE ARE NOT AN EXHAUSTIVE LIST OF METHODS OF PROOF. THEY ARE JUST THE MOST COMMON METHODS AND IT IS HOPED THAT THEY WILL SHOW HOW THE IDEAS OF MATHEMATICAL LOGIC CAN BE APPLIED IN STATING AND PROVING THEOREMS.



**Exercise 7.3**

- 1 PROVE THAT THE SUM OF TWO CONSECUTIVE ODD INTEGERS IS AN EVEN INTEGER.
- 2 SHOW THAT  $\sqrt{2}$  AND  $\sqrt{3}$  ARE IRRATIONAL NUMBERS, WHEN THERE EXISTS A RATIONAL NUMBER  $c$  SUCH THAT  $c < b$ .
- 3 PROVE THAT FOR ANY REAL NUMBERS  $a \geq 40$ , IF AND ONLY IF  $a \geq 20$  OR  $b \geq 20$ .
- 4 PROVE THAT THE SQUARE OF ANY INTEGER IS OF THE FORM  $4k$  OR  $4k + 1$ .
- 5 IF  $m, n \in \mathbb{N}$  AND  $m$  IS NOT A PERFECT SQUARE, THEN  $m$  PERFECT SQUARE OR NOT A PERFECT SQUARE IS A PERFECT SQUARE, IS SUCH THAT  $n^2$ .
- 6 SHOW THAT  $\sqrt{2}$  IS IRRATIONAL.
- 7 SHOW THAT  $\sqrt{x}$  AND  $\sqrt{y}$  ARE POSITIVE, THEN  $\sqrt{x^2 + y^2} \neq x + y$ .
- 8 CHECK WHETHER OR NOT EACH OF THE FOLLOWING IS TRUE.
  - A FOR ANY SETS  $A$  AND  $B \subseteq A \cup B$
  - B FOR ANY  $\mathbb{N}$ ,  $n$  IS EVEN IMPLIES THAT  $n$  IS NOT PRIME.
- 9 PROVE OR DISPROVE EACH OF THE FOLLOWING STATEMENTS
  - A IF  $x$  AND  $y$  ARE EVEN INTEGERS, IS  $x + y$  ALSO EVEN.
  - B IF  $3n + 2$  IS ODD, THEN  $n$  IS ODD.
  - C  $\forall n \in \mathbb{N}, n! < n^3$
  - D  $\forall n \in \mathbb{N}, n^2 < n^3$

**7.3 PRINCIPLE AND APPLICATION OF MATHEMATICAL INDUCTION**

BEFORE WE START WITH THE PRINCIPLE OF MATHEMATICAL INDUCTION, LET US CONSIDER SOME EXAMPLES.

**Example 1** CONSIDER THE SUM OF THE FIRST  $n$  ODD POSITIVE INTEGERS

IF $n = 1,$	$1 = 1$	$= 1^2$
IF $n = 2,$	$1 + 3 = 4$	$= 2^2$
IF $n = 3,$	$1 + 3 + 5 = 9$	$= 3^2$
IF $n = 4,$	$1 + 3 + 5 + 7 = 16$	$= 4^2$
IF $n = 5,$	$1 + 3 + 5 + 7 + 9 = 25$	$= 5^2$
IF $n = 6,$	$1 + 3 + 5 + 7 + 9 + 11 = 36$	$= 6^2$

FROM THE RESULTS ABOVE, IT LOOKS AS IF THE SUM OF THE FIRST  $n$  NATURAL NUMBERS IS ALWAYS GIVEN BY  $n^2$ . TO EXPRESS THIS IDEA SYMBOLICALLY, FIRST OBSERVE THE NATURAL NUMBER IS GIVEN BY  $n$  (WHICH YOU MAY CHECK YOURSELF). THEN WHAT WE HAVE DERIVED ABOVE CAN BE EXPRESSED AS:

$$1 + 3 + 5 + 7 + 9 + \dots + (2n - 1) = n^2 \quad (*)$$

YOU HAVE SEEN BY DIRECT CALCULATION THAT THE FORMULA (\*) IS TRUE WHEN  $n$  HAS THE VALUES 1, 2, 3, 4, 5 AND 6.

DOES THIS MEAN THAT THE FORMULA (\*) IS TRUE FOR ANY NATURAL NUMBER  $n$ ? CAN WE PROVE THIS SIMPLY BY CONTINUING NUMERICAL CALCULATIONS?

TRY THE CASE WHEN  $n = 13$ . DIRECT CALCULATION SHOWS THAT:

$$1 + 3 + 5 + 7 + 9 + 11 + 13 + 15 + 17 + 19 + 21 + 23 + 25 = 169 = 13^2.$$

SO, OUR FORMULA (\*) SEEMS TO HOLD. ONE MIGHT ALSO BE TEMPTED TO SAY THAT SINCE  $n$  IS CHOSEN RANDOMLY, THIS PROVES THAT (\*) IS TRUE FOR EVERY POSSIBLE CHOICE OF  $n$ . ACTUALLY, NO MATTER HOW MANY CASES YOU CHECK, YOU CAN NEVER PROVE (\*) IS ALWAYS TRUE, BECAUSE THERE ARE INFINITELY MANY CASES AND NO AMOUNT OF CALCULATION CAN CHECK THEM ALL.

SO, WHAT IS NEEDED IS SOME LOGICAL ARGUMENT THAT WILL PROVE THAT FORMULA (\*) IS TRUE FOR EVERY NATURAL NUMBER  $n$ .

BEFORE YOU CONSIDER THE DETAILS OF THIS LOGICAL ARGUMENT, SOME EXAMPLES OF STATEMENTS WHICH CAN BE CHECKED BY DIRECT CALCULATION FOR SOME VALUES OF  $n$  BUT WHICH TURN OUT TO BE FALSE FOR SOME OTHER VALUES OF  $n$ .

**Example 2** CONSIDER THE NUMBER  $P$  WHICH IS EXPRESSED IN THE FORM

$$P = 2^{2^n} + 1$$

WHERE  $n$  IS A NON-NEGATIVE INTEGER, THEN BY DIRECT CALCULATION, WE OBSERVE THAT

WHEN $n = 0$ ,	$P = 2^1 + 1 = 3$
WHEN $n = 1$ ,	$P = 2^2 + 1 = 5$
WHEN $n = 2$ ,	$P = 2^4 + 1 = 17$
WHEN $n = 3$ ,	$P = 2^8 + 1 = 257$
WHEN $n = 4$ ,	$P = 2^{16} + 1 = 65,537$

EACH OF THESE VALUES OF  $P$  IS A PRIME NUMBER. BASED ON THESE RESULTS, CAN YOU CONCLUDE THAT  $P$  IS ALWAYS A PRIME NUMBER FOR EVERY  $n$ ? OF COURSE NOT. YOU MIGHT GUESS THAT THIS IS TRUE BUT WE SHOULD NOT MAKE A POSITIVE STATEMENT UNLESS YOU CAN SUPPLY A PROOF THAT IS VALID FOR EVERY NATURAL NUMBER  $n$ . WHEN  $n = 5$ , THE NUMBER  $P$  IS FOUND NOT TO BE PRIME SINCE:

$$P = 2^{32} + 1 = 4,294,967,297 = 641 \times 6,700,417, \text{ WHICH IS NOT PRIME.}$$

**Example 3** CONSIDER THE INEQUALITY BELOW FOR A NATURAL NUMBER.

$$2^n < n^{10} + 2 \quad \dots\dots\dots \text{II}$$

IF WE CALCULATE BOTH SIDES OF (II) FOR THE FIRST FOUR VALUES OF  $n$

WHEN $n = 1$ , YOU GET	$2 < 1 + 2 = 3$
WHEN $n = 2$ , YOU GET	$4 < 1024 + 2 = 1026$
WHEN $n = 3$ , YOU GET	$8 < 59,051$
WHEN $n = 4$ , YOU GET	$16 < 1,048,578$

IT CERTAINLY APPEARS AS IF THE INEQUALITY IS TRUE FOR A NATURAL NUMBER. TRY FOR A LARGER VALUE OF  $n$ , THEN THE INEQUALITY SHOWS THAT


$$1,048,576 < 10,240,000,000,002$$

WHICH IS OBVIOUSLY TRUE. BUT, EVEN THIS DOES NOT PROVE THAT THE INEQUALITY IS TRUE. THIS ASSERTION IS ACTUALLY FALSE, BECAUSE WHEN APPROXIMATELY THAT  $2^{59} = 5.764 \times 10^{17}$  WHILE  $59^{10} + 2 = 5.111 \times 10^{17}$

THE LAST TWO EXAMPLES SHOW THAT YOU CANNOT CONCLUDE THAT AN ASSERTION INVOLVING INTEGERS IS TRUE FOR ALL POSITIVE VALUES BY CHECKING SPECIFIC VALUES OF  $n$ . NO MATTER HOW MANY YOU CHECK

HOW THEN IS SUCH AN ASSERTION PROVED TO BE TRUE?


AN ASSERTION INVOLVING A NATURAL NUMBER CAN BE PROVED BY USING A METHOD KNOWN AS THE PRINCIPLE OF MATHEMATICAL INDUCTION, STATED AS FOLLOWS.



## HISTORICAL NOTE

**Augustus Demorgan (1806 - 1871)**

One of the techniques to prove mathematical statements discussed in this unit is the Principle of Mathematical Induction. Even though the method was used by Fermat, Pascal and others before him, the actual term mathematical induction was first used by Demorgan. The method is used in many branches of higher mathematics.



**Principle of Mathematical Induction**

FOR A GIVEN ASSERTION INVOLVING A NATURAL NUMBER

- I THE ASSERTION IS TRUE FOR  $n = 1$
- II IT IS TRUE FOR  $n + 1$ , WHENEVER IT IS TRUE FOR  $n$ ,

THEN THE ASSERTION IS TRUE FOR EVERY NATURAL NUMBER  $n$

LET US NOW ILLUSTRATE THE USE OF THIS PRINCIPLE BY CONSIDERING DIFFERENT EXAMPLES. THE FIRST EXAMPLE WILL BE THE ONE WHICH YOU CONSIDERED AT THE BEGINNING OF THIS SECTION.

**Example 4** SHOW THAT THE SUM OF THE FIRST  $n$  NATURAL NUMBERS IS GIVEN BY  
 SHOW THAT,

$$1 + 3 + 5 + \dots + (2n - 1) = n^2 \dots\dots\dots *$$

FOR EVERY NATURAL NUMBER  $n$

**Proof:**

**1** IT IS CLEAR THAT THIS IS TRUE WHEN  $n = 1$  BECAUSE  $1^2 = 1$

**2** NOW ASSUME THAT THIS IS TRUE FOR  $k$ ; THAT IS ASSUME THAT

$$1 + 3 + 5 + \dots + (2k - 1) = k^2 \dots\dots\dots **$$

TO OBTAIN THE SUM OF THE FIRST  $k + 1$  ODD INTEGERS, YOU SIMPLY ADD THE NEXT ODD INTEGER WHICH IS  $2k + 1$  TO BOTH SIDES OF \*\* TO GET:

$$1 + 3 + 5 + \dots + (2k - 1) + (2k + 1) = k^2 + (2k + 1) = (k + 1)^2$$

THIS IS THE SAME AS REPLACING  $k$  WITH  $k + 1$ . HENCE, YOU HAVE SHOWN THAT IF THE ASSERTION IS TRUE FOR  $k$ , IT IS ALSO TRUE FOR  $k + 1$ .

BY THE PRINCIPLE OF MATHEMATICAL INDUCTION, THIS COMPLETES THE PROOF THAT THE ASSERTION IS TRUE FOR ANY NATURAL NUMBER  $n$ .

**Example 5** SHOW THAT THE EQUATION

$$1 + 4 + 7 + 10 + \dots + (3n - 2) = \frac{n(3n - 1)}{2} \dots\dots\dots I$$

IS TRUE FOR ANY NATURAL NUMBER  $n$

**Proof:**

**1** THE EQUATION IS TRUE FOR  $n = 1$  BECAUSE  $\frac{1(3(1) - 1)}{2} = \frac{1 \times 2}{2}$

**2** ASSUME THAT THE EQUATION IS TRUE FOR  $k$ ; THAT IS YOU ASSUME THAT,

$$1 + 4 + 7 + 10 + \dots + (3k - 2) = \frac{k(3k - 1)}{2} \dots\dots\dots II$$

NOW, IF YOU ADD THE NEXT ADDENDUM WHICH IS  $3k + 1$  TO BOTH SIDES OF II YOU GET:

$$\begin{aligned} 1 + 4 + 7 + 10 + \dots + (3k - 2) + (3k + 1) &= \frac{k(3k - 1)}{2} + (3k + 1) \\ &= \frac{k(3k - 1) + 2(3k + 1)}{2} = \frac{3k^2 + 5k + 2}{2} = \frac{(k + 1)(3k + 2)}{2} = \frac{(k + 1)(3(k + 1) - 1)}{2} \end{aligned}$$

BUT THIS LAST EQUATION IS THE SAME AS THE PREVIOUS ONE REPLACED BY  $n$ . HENCE YOU HAVE SHOWN THAT IF THE EQUATION IS TRUE FOR  $n$ , IT IS ALSO TRUE FOR  $n+1$  BY THE PRINCIPLE OF MATHEMATICAL INDUCTION. THIS COMPLETES THE PROOF THAT THE EQUATION IS TRUE FOR ANY NATURAL NUMBER.

**Example 6** PROVE THAT FOR ANY NATURAL NUMBER  $n$ ,  $1 < 2^n$ .

**Proof:**

- 1 FIRST FOR  $n=1$ ,  $1 < 2^1 = 2$  IS TRUE
- 2 ASSUME THAT  $1 < 2^n$  IS TRUE FOR  $n$ .

NOW YOU NEED TO SHOW IT IS TRUE ALSO FOR  $n+1$ .  $1 < 2^{n+1}$  IS ALSO TRUE.

ADDING 1 ON BOTH SIDES, YOU GET

$$n + 1 < 2^n + 1$$

AGAIN BECAUSE  $n$  IS FOR ANY NON-NEGATIVE INTEGER,  $n < 2^n$ .

$$n + 1 < 2^n + 1 \leq 2^n + 2^n = 2(2^n) = 2^{n+1}.$$

THUS  $n + 1 < 2^{n+1}$

THAT MEANS WHENEVER  $1 < 2^n$  IS TRUE,  $1 < 2^{n+1}$  IS ALSO TRUE. IN OTHER WORDS, WHENEVER YOUR ASSERTION IS TRUE FOR A NATURAL NUMBER  $n$ , IT IS ALSO TRUE FOR  $n+1$ .

THEREFORE, BY THE PRINCIPLE OF MATHEMATICAL INDUCTION, THE ASSERTION IS TRUE FOR ANY NATURAL NUMBER  $n$ .

**Example 7** USE MATHEMATICAL INDUCTION TO PROVE THAT  $n^3$  IS DIVISIBLE BY 3.

**Proof:**

- 1 THE ASSERTION IS TRUE WHEN  $n=1$  BECAUSE  $1^3 = 1$  AND 1 IS DIVISIBLE BY 3.
- 2 FOR  $n = k \geq 1$ , ASSUME THAT  $k^3$  IS DIVISIBLE BY 3 IS TRUE FOR A NATURAL NUMBER  $k$  AND YOU MUST SHOW THAT THIS IS ALSO TRUE FOR  $k+1$ . THIS MEANS YOU HAVE TO SHOW THAT  $(k+1)^3$  IS DIVISIBLE BY 3.

NOW, OBSERVE THAT

$$\begin{aligned} (k+1)^3 - k^3 &= (k^3 + 3k^2 + 3k + 1) - k^3 \quad (\text{EXPANDING } (k+1)^3) \\ &= (3k^2 + 3k + 1) \end{aligned}$$

SINCE BY THE ASSUMPTION  $k^3$  IS DIVISIBLE BY 3 AND  $3k^2$  IS CLEARLY DIVISIBLE BY 3, (AS IT IS 3 TIMES SOME INTEGER), YOU NOTICE THAT THE SUM  $3k^2 + 3k + 1$  IS DIVISIBLE BY 3. THUS, IT FOLLOWS THAT  $(k+1)^3$  IS DIVISIBLE BY 3. THEREFORE, BY THE PRINCIPLE OF MATHEMATICAL INDUCTION,  $n^3$  IS DIVISIBLE BY 3 FOR ANY NATURAL NUMBER  $n$ .

**Exercise 7.4**

1 SHOW THAT  $1 + 2 + 3 + \dots + n = \frac{n(n+1)}{2}$ , FOR EACH NATURAL NUMBER  $n$

2 SHOW THAT  $2 + 4 + 6 + \dots + 2n = 2n(n+1)$  FOR EACH NATURAL NUMBER  $n$

3 FIND  $2 + 4 + 6 + \dots + 100$ .

4 YOU MAY NOW ANSWER QUESTIONS 1 AND 2 OF THE OPENING PROBLEM OF THIS UNIT. PLEASE TRY THEM.

5 A SET OF BOXES ARE PUT ON TOP OF EACH OTHER. THE TOP ROW HAS 6 BOXES, THE ONE BELOW IT HAS 8 BOXES, AND THE NEXT LOWER ROWS HAS 10 BOXES AND SO ON. IF THERE ARE 41 ROWS AND 4110 BOXES ALL IN ALL, FIND THE VALUE OF  $n$

6 PROVE THAT THE  $n$ TH EVEN NATURAL NUMBER IS GIVEN BY  $2n$

7 PROVE THAT THE  $n$ TH ODD NATURAL NUMBER IS GIVEN BY  $2n - 1$

8 SHOW THAT 6 IS A MULTIPLE OF 5

9 SHOW THAT  $1 \leq n! \forall n \in \mathbb{N}$

10 SHOW THAT FOR ALL  $n \in \mathbb{N}$ ,  $1^3 + 2^3 + \dots + n^3 = \frac{n^2(n+1)^2}{4}$



**Key Terms**

- argument
- bi-implication
- conclusion
- conjunction
- connective
- counter example
- direct proof
- disjunction
- existential quantifier
- implication
- indirect proof (contra positive)
- mathematical induction
- method of cases (exhaustion)
- negation
- open statement
- premise
- proof by contradiction
- rules of inference
- statement (proposition)
- universal quantifier
- validity



## Summary

### 1 RULES OF CONNECTIVES: FOR PROPOSITIONS

$p$	$q$	$\neg p$	$p \wedge q$	$p \vee q$	$p \Rightarrow q$	$p \Leftrightarrow q$
T	T	F	T	T	T	T
T	F	F	F	T	F	F
F	T	T	F	T	T	F
F	F	T	F	F	T	T

### 2 **Universal quantifier:**

$\forall x$  MEANS FOR EVERY, FOR ANY, FOR EVERY, FOR ALL

### 3 **Existential quantifier:**

$\exists x$  MEANS FOR SOME THERE EXISTS

4  $(\forall x)(P(x) \Rightarrow Q(x))$ : EVERY  $x$  IS  $x$

5  $(\exists x)(P(x) \wedge Q(x))$ : SOME  $x$  IS  $x$  AND SOME  $(P)$  IS

6  $\neg(\forall x)P(x) \equiv (\exists x)\neg P(x)$

7  $\neg(\exists x)P(x) \equiv (\forall x)\neg P(x)$

8 AN ARGUMENT IS AN ASSERTION THAT A GIVEN SET OF STATEMENTS YIELD ANOTHER STATEMENT CALLED A

9 AN ARGUMENT IS VALID, IF WHENEVER ALL THE PREMISES ARE TRUE, THEN THE CONCLUSION IS TRUE. OTHERWISE IT IS CALLED A

10 AN ARGUMENT IS VALID, IF AND ONLY IF THE CONJUNCTION OF ALL PREMISES ALWAYS IMPLIES THE CONCLUSION.

### 11 **Rules of Inference:**

A  $\frac{p}{p \vee q}$  (Addition)

$p$

C  $\frac{q}{p \wedge q}$  (conjunction)

$p \Rightarrow q$

E  $\frac{\neg q}{\neg p}$  (Modus Tollens)

B  $\frac{p \wedge q}{p}$  (Simplification)

$p \Rightarrow q$

D  $\frac{p}{q}$  (Modus ponens)

$p \Rightarrow q$

F  $\frac{q \Rightarrow r}{p \Rightarrow r}$  (syllogism)

$$p \vee q$$

**G**  $\frac{\neg p}{q}$  (disjunctive syllogism)

**12 Direct proof:**

GIVEN A STATEMENT OF THE FORM  $P \Rightarrow Q$ , PROVING IT USING STEPS

$$\begin{array}{l} p \\ p_1 \\ p_2 \\ \vdots \\ p_n \\ \hline q \end{array}$$

WHERE  $p_1, p_2 \dots p_n$  ARE PREVIOUSLY ESTABLISHED THEOREMS OR IDENTITIES,  $P$  ETC, IS CALLED A **proof**.

**13 Method of cases:**

WHEN ONE PROVES AN ASSERTION BY CONSIDERING ALL POSSIBLE CASES, THE PROOF IS DONE BY METHOD OF **(cases)**.

**14 Indirect (contra positive) proof**

TO PROVE  $P \Rightarrow Q$  YOU CAN PROVE ITS CONTRA-POSITIVE

**15 Proof by contradiction**

TO SHOW THAT  $P$  IS TRUE, YOU SEEK FOR AN ASSERTION  $P \Rightarrow (r \wedge \neg r)$  IS TRUE.

**16 Disproving by counter example**

TO SHOW THAT  $P(x)$  IS FALSE, YOU SEEK AN OBJECT FROM THE UNIVERSE OF  $P(x)$  SUCH THAT IT IS FALSE (CALLED A **counter example**).

**17 Principle of mathematical induction**

IF FOR A GIVEN ASSERTION INVOLVING A NATURAL NUMBER  $n$ , YOU CAN SHOW THAT

- I THE ASSERTION IS TRUE FOR  $n = 1$
- II IF IT IS TRUE FOR  $n = k$  THEN IT IS ALSO TRUE FOR  $n = k + 1$

THEN THE ASSERTION IS TRUE FOR EVERY NATURAL NUMBER  $n$ .



**Review Exercises on Unit 7**

- 1** USING TRUTH TABLES, SHOW THAT EACH PAIR OF COMPOUND STATEMENTS ARE EQUIVALENT.
- A**  $p \Rightarrow q ; \neg p \vee q$                       **B**  $p \Leftrightarrow Q ; (p \Rightarrow q) \wedge (q \Rightarrow p)$   
**C**  $\neg (p \wedge q) ; \neg p \vee \neg q$                       **D**  $p \wedge (q \vee r) ; (p \wedge q) \vee (p \wedge r)$
- 2** USING TRUTH TABLES, SHOW THAT EACH OF THE FOLLOWING
- A**  $(p \wedge q) \Rightarrow p$                       **B**  $p \Rightarrow (p \vee q)$   
**C**  $[\neg p \wedge (p \vee q)] \Rightarrow q$                       **D**  $[p \wedge (p \Rightarrow q)] \Rightarrow q$
- 3** USE QUANTIFIERS TO EXPRESS EACH OF THE FOLLOWING STATEMENTS
- A** THERE IS A STUDENT IN THIS CLASS WHO CAN SPEAK FRENCH  
**B** EVERY STUDENT IN THIS CLASS KNOWS HOW TO DRIVE A CAR  
**C** THERE IS A STUDENT IN THIS CLASS WHO HAS A BICYCLE.
- 4** LET  $Q(x, y)$  BE THE OPEN PROPOSITION " $x^2 - y^2$ ". IF THE UNIVERSAL SET IS THE SET OF INTEGERS, WHAT ARE THE TRUTH VALUES OF THE FOLLOWING?
- A**  $Q(2, 2)$                       **B**  $Q(3, 0)$                       **C**  $(\forall x) Q(x, 0)$   
**D**  $(\exists x) Q(x, 4)$                       **E**  $(\exists x) (\exists y) Q(x, y)$                       **F**  $(\forall x) (\forall y) Q(x, y)$
- 5** IF, THE UNIVERSAL SET IS THE SET OF INTEGERS, DETERMINE THE TRUTH VALUE OF EACH OF THE FOLLOWING.
- A**  $(\forall n) (n^2 \geq 0)$                       **B**  $(\exists n) (n^2 = 2)$   
**C**  $(\forall n) (n^2 \geq n)$                       **D**  $(\forall n) (\exists m) (n < m^2)$   
**E**  $(\forall n) (\exists m) (n + m = 0)$                       **F**  $(\exists n) (\forall m) (nm = m)$   
**G**  $(\exists n) (\exists m) (n^2 + m^2 = 9)$                       **H**  $(\exists n) (\exists m) (n + m = 6 \wedge n - m = 2)$
- 6** IF, THE UNIVERSAL SET IS THE SET OF ALL REAL NUMBERS, DETERMINE THE TRUTH VALUE OF EACH OF THE FOLLOWING PROPOSITIONS.
- A**  $(\exists x) (x^2 = 3)$                       **B**  $(\exists x) (x^2 = -2)$   
**C**  $(\forall x) (\exists y) (x^2 = y)$                       **D**  $(\forall x) (\forall y) (x = y^2)$   
**E**  $(\exists x) (\forall y) (xy = 0)$                       **F**  $(\exists x) (\exists y) (xy \neq yx)$   
**G**  $(\forall x) (x \neq 0) (\exists y) (xy = 1)$
- 7** CHECK THE VALIDITY OF EACH OF THE FOLLOWING ARGUMENTS.
- |          |                                      |                        |   |
|----------|--------------------------------------|------------------------|---|
|          | $q \Rightarrow p$                    | $p \Rightarrow \neg q$ | $p \Rightarrow q$                             |
| <b>A</b> | $\frac{\neg q \Leftrightarrow p}{p}$ | <b>B</b>               | $\frac{p \wedge r}{\neg q \Leftrightarrow r}$ |
|          |                                      | <b>C</b>               | $\frac{p \Rightarrow r}{q \Rightarrow r}$     |

	$p \Rightarrow q$		$p \Rightarrow q$		$p \vee q$
<b>D</b>	$\frac{\neg p}{\neg q}$	<b>E</b>	$\frac{r \Rightarrow q}{p \Rightarrow r}$	<b>F</b>	$\frac{p}{\neg q}$

**8** CHECK THE VALIDITY OF EACH OF THE FOLLOWING ARGUMENTS.

**A** IF YOU SEND ME AN EMAIL MESSAGE, THEN I WILL GO TO SLEEP EARLY.  
 IF YOU DO NOT SEND ME AN EMAIL MESSAGE, THEN I WILL GO TO SLEEP EARLY.  
 IF I GO TO SLEEP EARLY, THEN I WILL WAKE UP EARLY.  
 THEREFORE, IF I DO NOT FINISH MY HOMEWORK, THEN I WILL WAKE UP EARLY.

**B** IF ALEMU HAS AN ELECTRIC CAR AND HE DRIVES A LONG DISTANCE, THEN HIS CAR WILL NEED TO BE RECHARGED. IF HIS CAR NEEDS TO BE RECHARGED, THEN HE WILL VISIT AN ELECTRIC STATION.  
 ALEMU DRIVES A LONG DISTANCE. HOWEVER, HE WILL NOT VISIT AN ELECTRIC STATION.  
 THEREFORE, ALEMU DOES NOT HAVE AN ELECTRIC CAR.

**9** PROVE OR DISPROVE EACH OF THE FOLLOWING STATEMENTS.

- A** IF  $x$  AND  $y$  ARE ODD INTEGERS, THEN  $x + y$  IS AN ODD INTEGER.
- B** THE PRODUCT OF TWO RATIONAL NUMBERS IS ALWAYS A RATIONAL NUMBER.
- C** THE PRODUCT OF TWO IRRATIONAL NUMBERS IS ALWAYS AN IRRATIONAL NUMBER.
- D** THE SUM OF TWO RATIONAL NUMBERS IS ALWAYS A RATIONAL NUMBER.
- E** IF  $n$  IS AN INTEGER AND IS ODD, THEN  $n^2$  IS EVEN.
- F** FOR EVERY PRIME NUMBER  $p$ ,  $p$  IS PRIME.
- G** FOR REAL NUMBERS  $p$  AND  $q$ , IF  $\sqrt{pq} \neq \frac{p+q}{2}$ , THEN  $p \neq q$ .
- H**  $\forall n, r \in \mathbb{Z}$  AND  $n \geq r \geq 2$ ,  $\binom{n}{r} = \binom{n}{n-r}$

**10** PROVE EACH OF THE FOLLOWING STATEMENTS BY THE METHOD OF MATHEMATICAL INDUCTION FOR ALL NATURAL NUMBERS

- A**  $1 + 2 + 2^2 + \dots + 2^n = \sum_{k=0}^n 2^k = 2^{n+1} - 1$
- B**  $1^2 + 2^2 + 3^2 + 4^2 + \dots + n^2 = \frac{n(n+1)(2n+1)}{6}$
- C**  $(1 \times 2) + (2 \times 3) + (3 \times 4) + \dots + n(n+1) = \frac{n(n+1)(n+2)}{3}$
- D**  $1^2 + 3^2 + 5^2 + \dots + (2n-1)^2 = \frac{n(2n-1)(2n+1)}{3}$
- E**  $\frac{1}{1 \times 2} + \frac{1}{2 \times 3} + \frac{1}{3 \times 4} + \dots + \frac{1}{n(n+1)} = \frac{n}{n+1}$